

APPLICATION
FOR
UNITED STATES LETTERS PATENT

TITLE: CARRIERS FOR CELL CULTURE AND METHODS FOR
CULTURING CELLS USING THE SAME

APPLICANT: MASAYUKI HARA, JUN MIYAKE AND AYAKO YAMAKI

CERTIFICATE OF MAILING BY EXPRESS MAIL

Express Mail Label No. EL558601271US

I hereby certify under 37 CFR §1.10 that this correspondence is being deposited with the United States Postal Service as Express Mail Post Office to Addressee with sufficient postage on the date indicated below and is addressed to the Commissioner for Patents, Washington, D.C. 20231.

October 27, 2000

Date of Deposit

Signature

Derek Norwood

Typed or Printed Name of Person Signing Certificate

00/2001-EE166960

2007 2008 2009 2010 2011 2012 2013 2014 2015 2016 2017 2018 2019 2020 2021 2022 2023 2024 2025 2026 2027 2028 2029 2030 2031 2032 2033 2034 2035 2036 2037 2038 2039 2040 2041 2042 2043 2044 2045 2046 2047 2048 2049 2050 2051 2052 2053 2054 2055 2056 2057 2058 2059 2060 2061 2062 2063 2064 2065 2066 2067 2068 2069 2070 2071 2072 2073 2074 2075 2076 2077 2078 2079 2080 2081 2082 2083 2084 2085 2086 2087 2088 2089 2090 2091 2092 2093 2094 2095 2096 2097 2098 2099 2100 2101 2102 2103 2104 2105 2106 2107 2108 2109 2110 2111 2112 2113 2114 2115 2116 2117 2118 2119 2120 2121 2122 2123 2124 2125 2126 2127 2128 2129 2130 2131 2132 2133 2134 2135 2136 2137 2138 2139 2140 2141 2142 2143 2144 2145 2146 2147 2148 2149 2150 2151 2152 2153 2154 2155 2156 2157 2158 2159 2160 2161 2162 2163 2164 2165 2166 2167 2168 2169 2170 2171 2172 2173 2174 2175 2176 2177 2178 2179 2180 2181 2182 2183 2184 2185 2186 2187 2188 2189 2190 2191 2192 2193 2194 2195 2196 2197 2198 2199 2200 2201 2202 2203 2204 2205 2206 2207 2208 2209 2210 2211 2212 2213 2214 2215 2216 2217 2218 2219 2220 2221 2222 2223 2224 2225 2226 2227 2228 2229 2230 2231 2232 2233 2234 2235 2236 2237 2238 2239 2240 2241 2242 2243 2244 2245 2246 2247 2248 2249 2250 2251 2252 2253 2254 2255 2256 2257 2258 2259 2260 2261 2262 2263 2264 2265 2266 2267 2268 2269 2270 2271 2272 2273 2274 2275 2276 2277 2278 2279 2280 2281 2282 2283 2284 2285 2286 2287 2288 2289 2290 2291 2292 2293 2294 2295 2296 2297 2298 2299 2300 2301 2302 2303 2304 2305 2306 2307 2308 2309 2310 2311 2312 2313 2314 2315 2316 2317 2318 2319 2320 2321 2322 2323 2324 2325 2326 2327 2328 2329 2330 2331 2332 2333 2334 2335 2336 2337 2338 2339 2340 2341 2342 2343 2344 2345 2346 2347 2348 2349 2350 2351 2352 2353 2354 2355 2356 2357 2358 2359 2360 2361 2362 2363 2364 2365 2366 2367 2368 2369 2370 2371 2372 2373 2374 2375 2376 2377 2378 2379 2380 2381 2382 2383 2384 2385 2386 2387 2388 2389 2390 2391 2392 2393 2394 2395 2396 2397 2398 2399 2400 2401 2402 2403 2404 2405 2406 2407 2408 2409 2410 2411 2412 2413 2414 2415 2416 2417 2418 2419 2420 2421 2422 2423 2424 2425 2426 2427 2428 2429 2430 2431 2432 2433 2434 2435 2436 2437 2438 2439 2440 2441 2442 2443 2444 2445 2446 2447 2448 2449 2450 2451 2452 2453 2454 2455 2456 2457 2458 2459 2460 2461 2462 2463 2464 2465 2466 2467 2468 2469 2470 2471 2472 2473 2474 2475 2476 2477 2478 2479 2480 2481 2482 2483 2484 2485 2486 2487 2488 2489 2490 2491 2492 2493 2494 2495 2496 2497 2498 2499 2500 2501 2502 2503 2504 2505 2506 2507 2508 2509 2510 2511 2512 2513 2514 2515 2516 2517 2518 2519 2520 2521 2522 2523 2524 2525 2526 2527 2528 2529 2530 2531 2532 2533 2534 2535 2536 2537 2538 2539 2540 2541 2542 2543 2544 2545 2546 2547 2548 2549 2550 2551 2552 2553 2554 2555 2556 2557 2558 2559 2560 2561 2562 2563 2564 2565 2566 2567 2568 2569 2570 2571 2572 2573 2574 2575 2576 2577 2578 2579 2580 2581 2582 2583 2584 2585 2586 2587 2588 2589 2590 2591 2592 2593 2594 2595 2596 2597 2598 2599 2600 2601 2602 2603 2604 2605 2606 2607 2608 2609 2610 2611 2612 2613 2614 2615 2616 2617 2618 2619 2620 2621 2622 2623 2624 2625 2626 2627 2628 2629 2630 2631 2632 2633 2634 2635 2636 2637 2638 2639 2640 2641 2642 2643 2644 2645 2646 2647 2648 2649 2650 2651 2652 2653 2654 2655 2656 2657 2658 2659 2660 2661 2662 2663 2664 2665 2666 2667 2668 2669 2670 2671 2672 2673 2674 2675 2676 2677 2678 2679 2680 2681 2682 2683 2684 2685 2686 2687 2688 2689 2690 2691 2692 2693 2694 2695 2696 2697 2698 2699 2700 2701 2702 2703 2704 2705 2706 2707 2708 2709 2710 2711 2712 2713 2714 2715 2716 2717 2718 2719 2720 2721 2722 2723 2724 2725 2726 2727 2728 2729 2730 2731 2732 2733 2734 2735 2736 2737 2738 2739 2740 2741 2742 2743 2744 2745 2746 2747 2748 2749 2750 2751 2752 2753 2754 2755 2756 2757 2758 2759 2760 2761 2762 2763 2764 2765 2766 2767 2768 2769 2770 2771 2772 2773 2774 2775 2776 2777 2778 2779 2780 2781 2782 2783 2784 2785 2786 2787 2788 2789 2790 2791 2792 2793 2794 2795 2796 2797 2798 2799 2800 2801 2802 2803 2804 2805 2806 2807 2808 2809 2810 2811 2812 2813 2814 2815 2816 2817 2818 2819 2820 2821 2822 2823 2824 2

FIELD OF THE INVENTION

BACKGROUND OF THE INVENTION

Among hydrated polymer gels, inter alia N-isopropylacrylamide (referred to as “NIPAM” hereafter), which is a temperature responsive polymer, can swell and is in a liquid state at a low temperature; however, it occurs phase transition at around 34°C thereby resulting in rapid contraction and gelation.

So far, for the purpose of stratification of cultured cell layers, a method has been used, wherein the cell cultured on NIPAM gel was piled up on another cell layer together with the NIPAM at a temperature of 37°C, subsequently the NIPAM was liquefied by lowering the temperature below 34°C to be removed, whereby those cells were directly piled up each other.

- 1 -

cells. In this case the cell must attach to ECM for its growth.

However, because the upper side of cell layer, as well as the region between the cell and NIPAM (as a basal layer), is not attached to another cell, ECM, which is necessary for cell adhesion, is not formed.

Thus, even if monolayers of cells cultured on NIPAM are piled up each other, and then the NIPAM is removed by solubilizing under temperature conditions below 34°C to pile up the cell layers so as to be contacted directly each other, the support is not enough for the cell overlaid above to be proliferated, accordingly, stable proliferation could not be expected.

Since the phenomenon is seen that the liquefied NIPAM also acts as cytotoxin to inhibit the normal cell growth, the above method was a very unsuitable and unstable technique as a means of cell stratification.

So far, there was no successful example of culturing a cell on a medium prepared by piling up the ECM component on various types of gels followed by gelation, and establishing a culture system has been attempted, the system employing a medium wherein the gel, no longer required after piling up, can be readily removed.

In those circumstances, the object of the present invention is to provide a technique for piling up cells in a stable and easy manner without using NIPAM which inhibits the cell growth and proliferation when it is solubilized, and a technique wherein cells on upper and lower sides can be adhered to each other via ECM (e.g. collagen) when the cell layers are piled up; that is, the invention is aimed at establishing a technique for the cell multi-layer which has been considered to be difficult to achieve *in vitro* except certain tissues including skin.

SUMMARY OF THE INVENTION

Alginic acid is a block copolymer composed of glucuronic acid (G) and mannuronic acid (M), wherein glucuronic acid forms an egg-box structure such that it surrounds a multivalent metal ion (e.g. calcium ion), thereby forming an alginate gel (see Fig.1). The greater the G/M ratio is, the higher the ability of alginic acid to form the gel becomes. However, since the alginate gel can not be molten even at a temperature above 100°C, the method using a

temperature sensitive substance like NIPAM is rather inappropriate as a method for exfoliating a living cell from alginate gel. On the other hand, the alginate gel is easily dissolved and liquefied when being soaked in a chelating agent (e.g. EDTA). Additionally, since the alginate gel is a natural product belonging to algae with a property of biodegradability, it does not inhibit the normal cell growth even when the alginate gel is not removed sufficiently upon its dissolution.

As a result of the extensive and intensive studies that were focused on the above properties of alginate gel, the present inventors have now found that it is easy to pile up a cell layer on an another cell layer when a cell is cultured using a carrier which comprises an alginate gel layer (e.g. calcium alginate gel layer) piled up on a porous membrane, followed by solubilizing the alginate gel layer. In this method utilizing the alginate gel, the cell culture containing cell layers can readily be detached.

The present inventors have also found that depending on types of cells to be cultured with the carrier for cell culture, the culture of cells can be carried out more efficiently by using a carrier for cell culture wherein ECM component gel layer (e.g. collagen gel layer) or ECM component sponge layer (e.g. collagen sponge layer) is further piled up on the alginate gel layer which has been piled up on a porous membrane.

The present invention was accomplished on the basis of the above described findings.

Thus, the present invention includes the following inventions:

- (1) A carrier for cell culture comprising a porous membrane and an alginate gel layer which is formed on the membrane.
- (2) The carrier of (1), wherein the alginate gel layer is composed of a calcium alginate gel.
- (3) The carrier of (1) or (2), which further comprises an extracellular matrix component gel layer or extracellular matrix component sponge layer which is formed on the alginate gel layer.
- (4) The carrier of (3), wherein the extracellular matrix component is collagen.
- (5) A method for culturing a cell, wherein the method comprises culturing the cell using the carrier for cell culture according to any one of (1) to (4).
- (6) A method for piling up a cell, wherein the method comprises: forming a cell layer on the

carrier according to any one of (1) to (4); solubilizing an alginate gel layer of the carrier thereby exfoliating the cell layer from a porous membrane of the carrier; and piling up the exfoliated cell layer on another cell layer formed on the carrier according to any one of (1) to (4).

(7) A cell multi-layer obtained by the method according to (6).

BRIEF DESCRIPTION OF THE DRAWING

FIG. 1 shows the procedures for preparing a carrier for cell culture.

FIG. 2 shows the procedures for culture of cells using the carrier for cell culture.

FIG. 3 shows the procedures for piling up cell layers.

DETAILED DESCRIPTION OF THE INVENTION

The present invention will be described in detail.

The carrier for cell culture of the invention is characterized by comprising a porous membrane and an alginate gel layer which is piled up on the membrane.

As used herein, the term "carrier for cell culture" means a carrier or support usable in culturing cells.

The term "porous membrane" as used herein means a membrane through which chelating agent is permeable but alginate gel is not. The "porous membrane" is not limited to particular ones so long as it satisfied the above definition, and it can include, in addition to a membrane with pores, a membrane with inner cavities or a membrane with both pores and inner cavities. Examples of the porous membrane includes a filter, an ultrafiltration membrane, a silicone rubber membrane, a polytetrafluoroethylene resin porous membrane (or PTFE porous membrane), a nonwoven fabric, a gauze-like mesh, and various types of membrane filters, preferably ultrafiltration membrane and hydrophilic PTFE porous membrane. Where the porous membrane has pores, the pore size is, but not limited to, generally $0.02-1000\ \mu\text{m}$, preferably $0.02-100\ \mu\text{m}$, more preferably $0.1-10\ \mu\text{m}$, so far as it is a size capable of permeating a chelating agent not an alginate gel.

As used herein, the term "alginate gel" means an alginate in which the gelation occurs due

Station	Time	Lat	Long	Alt	Temp	Hum	Wind	Dir	Speed	Pressure	Clouds	Remarks
1	0000	34° 15' N	118° 15' W	10	55.0	85	10	000	0.0	1013.2	000	Clear
2	0100	34° 15' N	118° 15' W	10	54.5	85	10	000	0.0	1013.2	000	Clear
3	0200	34° 15' N	118° 15' W	10	54.0	85	10	000	0.0	1013.2	000	Clear
4	0300	34° 15' N	118° 15' W	10	53.5	85	10	000	0.0	1013.2	000	Clear
5	0400	34° 15' N	118° 15' W	10	53.0	85	10	000	0.0	1013.2	000	Clear
6	0500	34° 15' N	118° 15' W	10	52.5	85	10	000	0.0	1013.2	000	Clear
7	0600	34° 15' N	118° 15' W	10	52.0	85	10	000	0.0	1013.2	000	Clear
8	0700	34° 15' N	118° 15' W	10	51.5	85	10	000	0.0	1013.2	000	Clear
9	0800	34° 15' N	118° 15' W	10	51.0	85	10	000	0.0	1013.2	000	Clear
10	0900	34° 15' N	118° 15' W	10	50.5	85	10	000	0.0	1013.2	000	Clear
11	1000	34° 15' N	118° 15' W	10	50.0	85	10	000	0.0	1013.2	000	Clear
12	1100	34° 15' N	118° 15' W	10	49.5	85	10	000	0.0	1013.2	000	Clear
13	1200	34° 15' N	118° 15' W	10	49.0	85	10	000	0.0	1013.2	000	Clear
14	1300	34° 15' N	118° 15' W	10	48.5	85	10	000	0.0	1013.2	000	Clear
15	1400	34° 15' N	118° 15' W	10	48.0	85	10	000	0.0	1013.2	000	Clear
16	1500	34° 15' N	118° 15' W	10	47.5	85	10	000	0.0	1013.2	000	Clear
17	1600	34° 15' N	118° 15' W	10	47.0	85	10	000	0.0	1013.2	000	Clear
18	1700	34° 15' N	118° 15' W	10	46.5	85	10	000	0.0	1013.2	000	Clear
19	1800	34° 15' N	118° 15' W	10	46.0	85	10	000	0.0	1013.2	000	Clear
20	1900	34° 15' N	118° 15' W	10	45.5	85	10	000	0.0	1013.2	000	Clear
21	2000	34° 15' N	118° 15' W	10	45.0	85	10	000	0.0	1013.2	000	Clear
22	2100	34° 15' N	118° 15' W	10	44.5	85	10	000	0.0	1013.2	000	Clear
23	2200	34° 15' N	118° 15' W	10	44.0	85	10	000	0.0	1013.2	000	Clear
24	2300	34° 15' N	118° 15' W	10	43.5	85	10	000	0.0	1013.2	000	Clear

Alginic acid exists naturally as a cell wall constitutive polysaccharide or intercellular filling material in phaeophyceae, from which alginic acid can be prepared. Example of phaeophyceae used as the raw material include Laminariales Laminaria (e.g. Laminaria japonica Areschoug). Alternatively, commercially available alginic acid may be used. The G/M ratio of alginic acid is not specifically limited to specific ones; however, the greater the G/M ratio is, the higher the ability of alginic acid to form a gel becomes. Thus, the G/M ratio is preferably high. More specifically, it is preferred to be 0.1 – 1, more preferably 0.2 – 0.5.

- 5 -

comprises a porous membrane and an alginate gel layer piled up thereon. For example, it can take the constitution wherein an ECM component gel layer or ECM component sponge layer is further piled up on the alginate gel layer. If a cell cultured using said carrier for cell culture is more apt to grow/develop on the ECM component gel layer or sponge layer than the alginate gel layer (e.g. fibroblast), it is preferred to pile up the ECM component gel layer or sponge layer on the alginate gel layer.

The term "extracellular matrix component (ECM) gel" as used herein means the gel matter of ECM component, and the term "ECM component gel layer" means a lamellar ECM component gel. The ECM is generally defined as "a stable bio-structure which exists outside the cell in animal tissue, being a complicated assembly of biopolymers extracellularly secreted and accumulated" (Dictionary of Biochemistry (3rd edition) p. 570, Tokyo Kagaku-dojin, Tokyo, Japan), and it plays roles in supporting cells materially and in regulating activity of cells (i.e. a role for the transduction of extracellular information to cells and the alteration of cellular activity). "Extracellular matrix component" as used herein means a component of extracellular matrix, such as collagen, elastin, proteoglycan, glucosaminoglycan, fibronectin, laminin, or vitronectin, preferably collagen or matrigel (which is a gel composed of collagen type IV, laminin, and heparan sulfate). The ECM components can be obtained by usual methods. Alternatively, commercially available ECM components may be used. The gelation of an ECM component can be carried out in the usual manner. For example, if the ECM component is collagen, then a collagen gel can be obtained by incubating an aqueous solution of 0.3-0.5% collagen at 37°C for 10-20 minutes. If necessary, a gelling agent may be used in gelation of an ECM component.

As used herein, the term "ECM component sponge" means an ECM component having been processed three-dimensionally into a porous sponge form, and the term "extracellular matrix component sponge layer" means a lamellar ECM component sponge. Since the ECM component sponge itself has a three-dimensional structure, the cell can be cultured in a pile up manner by using the ECM component sponge layer. Impregnating a liposome, which encapsulates various cell growth factors or growth factors therein, into inner vacancies of the ECM component sponge layer also enables cells present within the ECM component

sponge layer to induce differentiation without restraint. As for the ECM component, the same examples as mentioned above can be given. The ECM component sponge can be prepared in the usual manner. Commercially available ECM component sponges can also be used.

When the ECM component gel layer is piled up on alginate gel layer, the alginate gel layer and the ECM component gel layer can be piled up after preparing them separately. However, it is preferable to perform gelation of an aqueous solution containing the ECM component after the aqueous solution was overlaid on the alginate gel layer. This is because it is difficult to exfoliate the ECM component gel layer from a container (e.g. dish or schale) in which the ECM component gel layer is formed since the physical strength of the ECM component gel layer is not enough to be detached.

When the ECM component sponge layer is overlaid on the alginate gel layer, it is recommended to pile up the alginate gel layer and ECM component sponge layer after preparing them separately.

The thickness of the porous membrane, alginate gel layer and ECM component gel layer, which constitute the carrier for cell culture of the invention, is not particularly limited, but the thickness of the porous membrane is usually 0.01-1 mm, preferably 0.01-0.1 mm, more preferably 0.05-1 mm, the thickness of the alginate gel layer is usually 0.1-3 mm, preferably 1-2 mm, more preferably 1 mm, the thickness of ECM component gel layer is usually 0.1-1 mm, preferably 0.2-0.5 mm, more preferably 0.4 mm, the thickness of the ECM component sponge layer is usually 0.1-2 mm, preferably 0.2-1 mm, more preferably 0.5 mm.

The size of the carrier for cell culture usable in the invention can be determined appropriately according to for example the number of cells to be cultured, and the carrier can also be molded in an appropriate size by using a scalpel or the like.

The carrier for cell culture in the invention can be used for culturing cells. The culture of cells can be carried out, for example, on alginate gel layer, on ECM component gel layer, or on/in ECM component sponge layer. In the use of the carrier for cell culture composed of a porous membrane and an alginate gel layer piled up on the membrane as the carrier for cell culture of the invention, the cell can be cultured on the alginate gel layer. In the use of the

carrier for cell culture further comprising an ECM component gel layer piled up on the alginate gel layer as the carrier for cell culture of the invention, the cell can be cultured on the ECM component gel layer. In the use of the carrier for cell culture further comprising an ECM component sponge layer piled up on the alginate gel layer as the carrier for cell culture of the invention, the cell can be cultured on/in the ECM component sponge layer. Examples of the cell which can be cultured include fibroblast, vascular endothelial cell, chondrocyte, hepatocyte, small intestine epitheliocyte, epidermis cornification cell, osteoblast, and bone marrow mesenchymal cell, preferably fibroblast. When cell is cultured, a culture medium (e.g. D-MEM, MEM, HamF12 or HamF10 medium) with a cell concentration of 10,000 – 15,000 cells/ml is usually added onto the alginate gel layer, ECM component gel layer, or ECM component sponge layer. The culture conditions of cells can be selected appropriately according to types of cells to be cultured. When cell is cultured on alginate gel layer or ECM component gel layer, the culture is usually continued until confluent cell monolayer is formed on the alginate gel layer or ECM component gel layer.

Specifically, the culture of cells using the carrier for cell culture of the invention can be carried out as follows. The carrier for cell culture is placed in, for example, a schale, and an appropriate culture medium (e.g. D-MEM, MEM, HamF12 or HamF10 medium) is added into the schale, then left for 12-24 hours to soak the culture medium into the carrier for cell culture. The culture medium in the schale is decanted, cells are spread on the alginate gel layer, ECM component gel layer or ECM component sponge layer, and subsequently an appropriate culture medium (e.g. D-MEM, MEM, HamF12 or HamF10 medium) is added into the schale. After leaving 1-2 hours at 37°C to attach the cell to the alginate gel layer, ECM component gel layer, or ECM component sponge layer, the cell culture is continued at 37°C. During the culture, the culture medium may be exchanged with the same fresh medium if necessary. It is usually exchanged in 0.5-2 day intervals after start of the culture.

The cell culture obtained by the culture of cells using the carrier for cell culture of the invention contains the carrier for cell culture of the invention and the cell layer maintained on the carrier. The "cell layer maintained by the carrier" contains any one of or combination of a cell layer formed on the alginate gel layer, a cell layer formed on the ECM component gel

layer, and a cell layer formed on/in the ECM component sponge layer. In the use of the carrier for cell culture composed of a porous membrane and an alginate gel layer piled up thereon as the carrier of the invention, or in the use of the carrier for cell culture further comprising an ECM component gel layer piled up on alginate gel layer as the carrier of the invention, or in the use of the carrier for cell culture further comprising an ECM component sponge layer piled up on alginate gel layer as the carrier of the invention, the cell layer formed in and/or on the ECM component sponge layer corresponds to the "cell layer maintained by the carrier."

The cell culture can be detached from the porous membrane by solubilizing the alginate gel layer. The solubilization of the alginate gel layer can be carried out by use of a chelating agent. Examples of the chelating agent include polyaminocarboxylic acids such as EDTA (ethylenediaminetetraacetic acid) and EGTA (ethylene glycol-bis(β -aminoethyl ether)); and oxycarboxylic acids such as citric acid, preferably EDTA and EGTA. The chelating agent can be selected appropriately according to the type of a multivalent metal ion which forms a chelate structure with a carboxylic acid group in the molecule of alginic acid. In the case of calcium alginate gel, for example, EDTA can be used, wherein the concentration of the chelating agent is usually 0.01-1 M, preferably 0.05-0.2 M. The solubilization of the alginate gel layer with a chelating agent is preferably carried out by allowing the chelating agent to soak through the porous membrane. Hereby, the porous membrane and alginate gel layer can be separated easily, and the cell culture can readily be detached from the porous membrane. It is not necessary to completely remove the alginate gel layer by solubilization, and the alginate gel layer incapable of being solubilized may remain. Preferably, the alginate gel layer is removed by solubilizing as much as possible.

Since the cell culture obtained by solubilizing the alginate gel layer contains a cell layer, it can be used for forming cell multi-layer. In forming cell multi-layer, the cell cultures obtained by solubilizing the alginate gel layer may be piled up each other, or one cell culture obtained by solubilizing the alginate gel layer may be piled up on another cell layer prepared separately. The type of cell of a cell layer to be piled up may be the same or different. The number of cell layers to be piled up is, but not limited to, usually 1-10, preferably 1-5, more

preferably 1-3.

The cell multi-layer contains any one of or combination of a cell layer formed on alginate gel layer, a cell layer formed on ECM component gel layer, and a cell layer formed on/in the ECM component sponge layer.

Used as the cell layer to be piled up are for example small intestine epitheliocyte layer, muscular layer, and fibroblast layer. By use of such cell layers, the three-dimensional tissue structure of the small intestine wall can be constructed. This three-dimensional tissue structure can be applied as an alternative model for animal experiments or as an internal organ for transplantation, as well as to an *in vitro* drug permeability test.

The cell multi-layer can be cultured under appropriate culture conditions depending on the cell type which composes a cell layer. In culture, a medium such as D-MEM, MEM, HamF12 or HamF10 medium can be used.

EXAMPLES

EXAMPLE 1: Preparation of Carrier for Cell Culture

The procedures for the preparation of a carrier for cell culture are shown in FIG. 1. Specifically, the carrier was prepared as follows;

- (1) One ml of a solution of 1% sodium alginate in water was added to a cell of which bottom is a porous membrane (pore size $3.0\mu\text{m}$; available from FALCON). The porous membrane positioned underneath can permeate water molecule and relatively small molecules, but it cannot permeate macromolecules, such as cultured cells, and polymer gels.
- (2) Gelation was performed by allowing 0.1 M CaCl_2 to soak through the porous membrane at the bottom of the cell and leaving for 1 hour at room temperature, thereby to prepare a carrier for cell culture that comprises a porous membrane and a calcium alginate gel layer piled up thereon.
- (3) 0.5 ml of a 3% collagen solution (collagen type IAC-50, available from Koken, Tokyo, Japan) diluted with D-MEM medium (available from Sigma) was added onto the calcium alginate gel layer to prepare a thin layer of the aqueous collagen solution,

followed by incubation for about 20 minutes at 37°C in a CO₂ incubator, thereby resulting in gelation of the collagen solution. By this means, the carrier for cell culture wherein the collagen gel layer was piled up on the calcium alginate gel layer, was obtained.

EXAMPLE 2 : Cell Culture using the Carrier for Cell Culture

The procedures for the culture of a cell using the carrier for cell culture are shown in FIG.2. Specifically, the cell culture was carried out as follows.

- (1) After placing the carrier for cell culture prepared in Example 1 along with the cell into a schale, D-MEM medium was added in the amount of 2ml into the cell and in the amount of 3ml into the schale, then left overnight in order to allow the D-MEM medium to soak into the carrier for cell culture,
- (2) Fibroblast cultured in advance was collected by trypsin treatment, and the cell concentration was adjusted to 20,000cells/ml. After removing the cell and the medium inside the schale, 0.5 ml of the cultured fibroblast (where the number of cells is 10,000) was added onto the collagen gel layer, and 3 ml of D-MEM medium was added into the schale,
- (3) After incubating for 1 hour at 37°C in a CO₂ incubator, the fibroblast was attached and retained on the collagen gel layer,
- (4) The medium was exchanged with the fresh one on Day 2 after culture, and the confluent cell monolayer was formed following culture of one more day. As a result, the cell culture which comprises a porous membrane, a calcium alginate gel layer piled up on the porous membrane, a collagen gel layer piled up on the calcium alginate gel layer, and a fibroblast layer formed on the collagen gel layer, was obtained (see FIG. 2).

EXAMPLE 3: Piling Up Cell Layer

The procedures for piling up a cell layer are shown in FIG.3. Specifically, they were performed as follows.

- (1) The cell culture obtained in Example 2 along with the cell was soaked in the 0.1M EDTA

solution, then EDTA was permeated through the porous membrane to dissolve the calcium alginate gel. By this means, the cell culture comprising the fibroblast layer could be detached from the porous membrane.

- (2) An excess of water was removed by suction, and a scalpel was inserted through the inner wall of the cell to hollow out the porous membrane, whereby the cell culture containing the fibroblast layer was suspended in D-MEM medium.
- (3) The cell culture containing this fibroblast was piled up on another cell culture obtained in the same manner as in Example 2. By repeating the same operations, piled up cell layers (consisting of 3 layers) were obtained.
- (4) It was confirmed that the piled up cell layers could be cultured in D-MEM medium.

As demonstrated by the above described examples, according to the present invention there is provided a carrier for cell culture capable of piling up cell layers easily.